

Annual PQRS Questionnaire (Pain)

Patient Name: _____ Date: _____

1. Do you have little or no interest in doing things? No
 Yes, please check one: Several Days More than half the days Everyday

2. Are you feeling down, depressed or hopeless? No
 Yes, please check one: Several Days More than half the days Everyday

If you answered YES to question 1 or 2, then complete the following table. If you answered No to both question 1 and 2 then you DO NOT have to complete the table, skip to Immunizations below.

	Not at all (0)	Several days (1)	More than half the days (2)	Everyday (3)
Do you have trouble falling or staying asleep or sleeping too much?				
Do you feel tired or have little energy?				
Do you have poor appetite or overeating?				
Do you feel bad about yourself, feel like a failure, or feel you have let yourself or your family down?				
Do you have trouble concentrating on things, such as reading the newspaper or watching television?				
Do you move or speak so slowly that other people could have noticed? Or the opposite? Are you fidgety or restless and move around more than usual?				
Do you have thoughts that you would be better off dead and/or have thoughts of hurting yourself in some way?				

Immunizations:

Have you had a Pneumonia Vaccination? NO YES, when: _____

Have you had a Flu Vaccination? No YES, when: _____

Have you fallen in the past year? (Answer only if 65 years and older.)

NO

Yes : 1 fall with injury 2 or more falls with injury
 1 fall without injury 2 or more falls without injury

Opioid Risk Tool

This tool should be administered to patients upon an initial visit prior to beginning opioid therapy for pain management. A score of 3 or lower indicates low risk for future opioid abuse, a score of 4 to 7 indicates moderate risk for opioid abuse, and a score of 8 or higher indicates a high risk for opioid abuse.

Mark each box that applies	Female	Male
Family history of substance abuse		
Alcohol	1	3
Illegal drugs	2	3
Rx drugs	4	4
Personal history of substance abuse		
Alcohol	3	3
Illegal drugs	4	4
Rx drugs	5	5
Age between 16—45 years	1	1
History of preadolescent sexual abuse	3	0
Psychological disease		
ADD, OCD, bipolar, schizophrenia	2	2
Depression	1	1
Scoring totals		

Questionnaire developed by Lynn R. Webster, MD to assess risk of opioid addiction.

Webster LR, Webster R. Predicting aberrant behaviors in Opioid-treated patients: preliminary validation of the Opioid risk tool. *Pain Med.* 2005; 6 (6) : 432

Oswestry Low Back Pain Disability Questionnaire

Instructions

This questionnaire has been designed to give us information as to how your back or leg pain is affecting your ability to manage in everyday life. Please answer by checking ONE box in each section for the statement which best applies to you. We realise you may consider that two or more statements in any one section apply but please just shade out the spot that indicates the statement which most clearly describes your problem.

Section 1 – Pain intensity

- I have no pain at the moment
- The pain is very mild at the moment
- The pain is moderate at the moment
- The pain is fairly severe at the moment
- The pain is very severe at the moment
- The pain is the worst imaginable at the moment

Section 2 – Personal care (washing, dressing etc)

- I can look after myself normally without causing extra pain
- I can look after myself normally but it causes extra pain
- It is painful to look after myself and I am slow and careful
- I need some help but manage most of my personal care
- I need help every day in most aspects of self-care
- I do not get dressed, I wash with difficulty and stay in bed

Section 3 – Lifting

- I can lift heavy weights without extra pain
- I can lift heavy weights but it gives extra pain
- Pain prevents me from lifting heavy weights off the floor, but I can manage if they are conveniently placed eg. on a table
- Pain prevents me from lifting heavy weights, but I can manage light to medium weights if they are conveniently positioned
- I can lift very light weights
- I cannot lift or carry anything at all

Section 4 – Walking*

- Pain does not prevent me walking any distance
- Pain prevents me from walking more than 1 mile
- Pain prevents me from walking more than 1/2 mile
- Pain prevents me from walking more than 100 yards
- I can only walk using a stick or crutches
- I am in bed most of the time

PATIENT NAME: _____

Date: _____

Section 5 – Sitting

- I can sit in any chair as long as I like
- I can only sit in my favourite chair as long as I like
- Pain prevents me sitting more than one hour
- Pain prevents me from sitting more than 30 minutes
- Pain prevents me from sitting more than 10 minutes
- Pain prevents me from sitting at all

Section 6 – Standing

- I can stand as long as I want without extra pain
- I can stand as long as I want but it gives me extra pain
- Pain prevents me from standing for more than 1 hour
- Pain prevents me from standing for more than 30 minutes
- Pain prevents me from standing for more than 10 minutes
- Pain prevents me from standing at all

Section 7 – Sleeping

- My sleep is never disturbed by pain
- My sleep is occasionally disturbed by pain
- Because of pain I have less than 6 hours sleep
- Because of pain I have less than 4 hours sleep
- Because of pain I have less than 2 hours sleep
- Pain prevents me from sleeping at all

Section 8 – Sex life (if applicable)

- My sex life is normal and causes no extra pain
- My sex life is normal but causes some extra pain
- My sex life is nearly normal but is very painful
- My sex life is severely restricted by pain
- My sex life is nearly absent because of pain
- Pain prevents any sex life at all

Section 9 – Social life

- My social life is normal and gives me no extra pain
- My social life is normal but increases the degree of pain
- Pain has no significant effect on my social life apart from limiting my more energetic interests eg, sport
- Pain has restricted my social life and I do not go out as often
- Pain has restricted my social life to my home
- I have no social life because of pain

Section 10 – Travelling

- I can travel anywhere without pain
- I can travel anywhere but it gives me extra pain
- Pain is bad but I manage journeys over two hours
- Pain restricts me to journeys of less than one hour
- Pain restricts me to short necessary journeys under 30 minutes
- Pain prevents me from travelling except to receive treatment

References

1. Fairbank JC, Pynsent PB. The Oswestry Disability Index. Spine 2000 Nov 15;25(22):2940-52; discussion 52.